

Alaska Department of Fish and Game

Division of Commercial Fisheries Professional Paper



Professional papers include published manuscripts that address management issues or research undertaken by the Alaska Department of Fish and Game, and are prepared by division personnel for publication in a professional journal. The primary audience will be fishery professionals, but some papers may be of interest to other disciplines. For access to these papers see the electronic links provided below.

Narum, S. R., M. Banks, T. D. Beacham, R. Bellinger, M. Campbell, J. DeKoning, A. Elz, C. M. Guthrie, C. Kozfkay, K. Miller, P. Moran, R. Phillips, L. W. Seeb, C. T. Smith, K. Warheit, S. Young, and J. C. Garza. 2008. Differentiating salmon populations at broad and fine geographic scales with microsatellites and single nucleotide polymorphisms. *Molecular Ecology* 17:3464-3477.

Abstract: Single nucleotide polymorphisms (SNPs) are appealing genetic markers due to several beneficial attributes, but uncertainty remains about how many of these bi-allelic markers are necessary to have sufficient power to differentiate populations, a task now generally accomplished with highly polymorphic microsatellite markers. In this study, we tested the utility of 37 SNPs and 13 microsatellites for differentiating 29 broadly distributed populations of Chinook salmon ($n = 2783$). Information content of all loci was determined by I_n and $G'ST$, and the top 12 markers ranked by I_n were microsatellites, but the 6 highest, and 7 of the top 10 $G'ST$ ranked markers, were SNPs. The mean ratio of random SNPs to random microsatellites ranged from 3.9 to 4.1, but this ratio was consistently reduced when only the most informative loci were included. Individual assignment test accuracy was higher for microsatellites (73.1%) than SNPs (66.6%), and pooling all 50 markers provided the highest accuracy (83.2%). When marker types were combined, as few as 15 of the top ranked loci provided higher assignment accuracy than either microsatellites or SNPs alone. Neighbour-joining dendrograms revealed similar clustering patterns and pairwise tests of population differentiation had nearly identical results with each suite of markers. Statistical tests and simulations indicated that closely related populations were better differentiated by microsatellites than SNPs. Our results indicate that both types of markers are likely to be useful in population genetics studies and that, in some cases, a combination of SNPs and microsatellites may be the most effective suite of loci.

Keywords: assignment tests, Chinook salmon, differentiation, genetic markers, microsatellites, SNPs

DOI: [10.1111/j.1365-294X.2008.03851.x](https://doi.org/10.1111/j.1365-294X.2008.03851.x)

You may be directed away from the ADF&G website. You may be required to pay a fee to view the full-text article. Library assistance may be available through the Alaska Resources Library and Information Services (ARLIS) <http://www.adfg.alaska.gov/index.cfm?adfg=libraryarlis.main> or through your local library.

Paper No. NA